ISSN (print) 0093-4666 © 2015. Mycotaxon, Ltd. ISSN (online) 2154-8889

MYCOTAXON

http://dx.doi.org/10.5248/130.223

Volume 130, pp. 223–225 *January–March 2015*

Hyphoderma hallenbergii, a new corticioid species from India

Maninder Kaur, Avneet P. Singh*, & G.S. Dhingra

Department of Botany, Punjabi University, Patiala 147002, India

*Correspondence to: *avneetbot@gmail.com*

Abstract — *Hyphoderma hallenbergii* sp. nov. from Himachal Pradesh, India is described.

Key words — Baghi, *Basidiomycota*, *Agaricomycetes*, *Meruliaceae*

During August 2012 fungal forays conducted of the Baghi area of district Shimla, Himachal Pradesh, India, Maninder and Avneet collected a fungus on decaying wood of *Cedrus deodara* in a mixed forest. Comparison of macroscopic and microscopic characters (Eriksson & Ryvarden 1975, Dhingra 1989, Larsson 1998, Bernicchia & Gorjón 2010, Singh et al. 2010, Priyanka & Dhingra 2012, Mycobank 2014) suggested that it belonged in the genus *Hyphoderma* (*Meruliaceae*, *Agaricomycetes*, *Basidiomycota*). The fungus, which is similar to *H. nemorale* K.H. Larss. and *H. incrustatum* K.H. Larss. in having two types of cystidia but which differs in basidiospore size, is described here as a new species.

Hyphoderma hallenbergii Man. Kaur, Avneet P. Singh & Dhingra, **sp. nov.** Plate 1

MycoBank MB 809801

Differs from *Hyphoderma nemorale* and *H. incrustatum* by its smaller and ellipsoid basidiospores.

Type: India, Himachal Pradesh: Shimla, Narkanda, on Baghi road, on the bark of decaying wood of *Cedrus deodara* (Roxb. ex D. Don) G. Don, 19 August 2012, Maninder Kaur & Avneet 6962 (PUN, **holotype**).

Etymology: In the honor of Dr. Nils Hallenberg, Professor Emeritus, University of Gothenburg, Sweden.

Basidiocarp resupinate, adnate, effused, ≤100 µm thick in section; hymenial surface smooth to somewhat grandinioid, chalky white; margins thinning



Plate 1. *Hyphoderma hallenbergii* (holotype). 1. Basidiocarp showing hymenial surface. 2. Capitate hymenial cystidia. 3. Basidium. 4,5. Basidiospores. 6. Vertical section through basidiocarp.

out or indeterminate, concolorous or paler. Hyphal system monomitic. Generative hyphae ≤2.5 µm wide, branched, septate, clamped; subiculum poorly developed, formed of few hyphae running more or less parallel to the substrate; subhymenial hyphae well developed, vertical, denser. Cystidia of two types: [1] Capitate cystidia 25–44 × 5–6 µm, subfusiform, originating from the hymenium, thin-walled, with basal clamp, sparsely encrusted; [2] Cystidia which are cylindrical to flexuous with constrictions, sometimes almost moniliform 60–80 × 5–7 µm, originating from subhymenium, slightly thick-walled, with basal clamp, often encrusted with crystals. Basidia 16–34 × 4–5 µm, clavate to subclavate, constricted to sinuous, 4-sterigmate, with basal clamp; sterigmata ≤3.5 µm long. Basidiospores 6.3–7.5 × 3.5–4.5 µm, ellipsoid, with oily contents, thin-walled, inamyloid, acyanophilous.

Remarks — *Hyphoderma nemorale* and *H. incrustatum* both differ from *H. hallenbergii* by their larger, cylindrical basidiospores (*H. nemorale*, 10–14 × 4–5 µm; *H. incrustatum*, 11–16 × 4–5.5 µm; Larsson 1998).

Acknowledgements

The authors thank Head (Department of Botany, Punjabi University, Patiala) for providing research facilities, Dr. Nils Hallenberg (Professor Emeritus, Denmark) for expert comments and peer review; and Prof. B.M. Sharma (Department of Plant Pathology, COA, CSKHPAU, Palampur, H.P., India) for peer review.

Literature cited

Bernicchia A, Gorjón SP. 2010. *Corticiaceae* s.l. Fungi Europaei 12. Edizioni Candusso. Alassio. Italia. 1008 p.

Dhingra GS. 1989. Genus *Hyphoderma* Wallr. em. Donk in the Eastern Himalayas. 197–212, in: ML Triveda et al. (eds). Plant Science Research in India. Today & Tomorrow's Printers & Publishers, New Delhi.

Eriksson J, Ryvarden L. 1975. The *Corticiaceae* of North Europe– III. Fungiflora, Oslo. pp. 288–546.

Larsson KH. 1998. Two new species in *Hyphoderma*. Nordic Journal of Botany. 18(1): 121–127. http://dx.doi.org/10.1111/j.1756-1051.1998.tb01106.x

MycoBank. 2014. Fungal databases. Nomenclature and species banks. [Accessed: 16/07/2014] http://www.mycobank.org/.

Priyanka, Dhingra GS. 2012. Two new species of *Hyphoderma* (*Agaricomycetes*) from India. Mycotaxon 119: 255–260. http://dx.doi.org/10.5248/119.255

Singh AP, Priyanka, Dhingra GS, Singla N. 2010. A new species of *Hyphoderma* (*Basidiomycetes*) from India. Mycotaxon 111: 71–74. http://dx.doi.org/10.5248/111.71